



Mountain West Gastroenterology

Colonoscopy Prep Instructions – COLYTE PREPARATION 2 DAY PREP

<input type="checkbox"/> Oscar Batista, M.D.	<input type="checkbox"/> Peter Fenton, M.D.	<input type="checkbox"/> Randall Ryser, M.D.
<input type="checkbox"/> Mark Boschert, M.D.	<input type="checkbox"/> Edward Frech, M.D.	<input type="checkbox"/> Michael Sossenheimer, M.D.
<input type="checkbox"/> John Bowers, M.D.	<input type="checkbox"/> Gordon Harmston, M.D.	<input type="checkbox"/> Raymond Thomason, M.D.
<input type="checkbox"/> Christopher Canale, M.D.	<input type="checkbox"/> Clarke Hilbig, M.D.	
<input type="checkbox"/> Brett Doxey, M.D.	<input type="checkbox"/> Robert Jones, M.D.	

7 DAYS PRIOR TO YOUR EXAM:

- Stop taking **Aspirin, Ginkgo Biloba, or Plavix**. Tylenol is ok to take. If you have a heart condition, had a recent heart attack or cardiac stent placement, you should discuss this with your Cardiologist or Primary Care Physician prior to stopping any anti-clotting agents.
- If you are taking blood thinners such as **Coumadin**, you should discuss with your Cardiologist or Primary Care Physician how to manage your blood thinners prior to the procedure.
- If you have had a mechanical heart valve, history of deep clots in your legs, or a pulmonary embolism, you may need **Lovenox or Heparin** up to the day of your procedure and possibly after the procedure.
- If you have **diabetes**, please follow our attached guidelines and discuss this with your Primary Care Physician. Test your blood sugar before coming in for your exam and bring the reading with you. If you take oral medications for your diabetes, don't take them the morning of your procedure unless otherwise instructed by your doctor. If you take insulin, bring your insulin with you. If you take insulin in the morning, take only ½ of your usual dose.
- **Nursing Mothers** – please “pump and dump” twice after the procedure before resuming breastfeeding.

5 DAYS PRIOR TO YOUR EXAM:

Avoid eating high fiber foods, fiber supplements and discontinue any **iron** supplements 5 days prior to your examination.

3 DAYS PRIOR TO YOUR EXAM:

Stop taking all anti-inflammatory medicines such as **Ibuprofen, Advil, Naproxen, Excedrin, Aleve or similar**. You can take Tylenol for pain.

Have your prescription for Colyte prep filled at your pharmacy.

Buy one 10 oz bottle of Magnesium Citrate.

2 DAYS PRIOR TO YOUR EXAM: *You may not eat any solid food until after your procedure!*

Drink only “Clear Liquids” for BREAKFAST, LUNCH and DINNER. This means liquids you can see through such as Apple juice, water, Sprite, Jell-O, Coffee (No Creamer), tea, clear chicken or beef broth, clear Gatorade, clear Powerade, or Crystal Light. *NO milk or pulp products: do not drink anything with red, purple or orange dyes. Drink as many liquids as you would like – keep yourself hydrated.

At **7:00 pm** drink on 10 oz. bottle of Magnesium Citrate. You may continue to drink clear liquids.

1 DAY PRIOR TO YOUR EXAM: *You may not eat any solid food until after your procedure!*

One of the most important aspects to a successful prep is to consume plenty of fluids the day before the exam.

In the morning, add water to fill the Colyte jug, shake well, and refrigerate. If you prefer, just add cold water to the container a couple hours before you start drinking it. Do not add anything to the Colyte preparation such as soda, etc.

Restrict your diet to clear liquids only, no solids. Liquids can be taken without restriction.

By **5:00 pm**, start drinking your Colyte prep. Drink ½ gallon of the Colyte (total 2 liters) over 2 hours.

You may continue to drink water to stay hydrated.

THE DAY OF YOUR EXAM: *You may not eat any solid food until after your procedure!*

- If you have diabetes, test your blood sugar before coming in for your exam and bring the reading with you. If you take oral medications for your diabetes, don't take them the morning of your procedure unless otherwise instructed by your doctor. If you take insulin, bring your insulin with you. If you take insulin in the morning, take only ½ of your usual dose.
- Remember to TAKE YOUR USUAL MEDICATIONS (no blood thinners, no NSAIDS, no aspirin, no Plavix) the morning of your procedure.
- At least 4-5 hours before your scheduled arrival time at the endoscopy center, drink the remaining ½ gallon of the Colyte (2 liters) over 2 hours. **Be sure to finish the bottle.** You may experience headache, chills, or nausea from the Colyte; do not be alarmed. You may have to sit on the toilet while taking the Colyte. When all the Colyte is gone, you may continue to drink water until two hours prior to your scheduled arrival time. Stop drinking 2 hours before your scheduled arrival time.

You may apply zinc oxide, Vaseline, Aloe Wipes or petroleum jelly to anal area to prevent irritation. Baby Wipes or Medicated Tucks may be considered.

Please bring someone with you to the endoscopy center to drive you home, as you will be sedated for the exam. You may not drive for 12 hours.

Your exam will be cancelled or must be done without sedation if you do not have someone to drive you home. You will be in the Endoscopy Lab for approximately two to three hours. The doctor will talk to you after the exam and will give you recommendations for diet, medication, follow up care, etc. Wear comfortable clothing; bring your glasses, hearing aids, insurance card(s) and completed information form. We will expect payment of co-pays, coinsurance and deductibles at the time of service. If your insurance requires a referral, you should contact your primary care physician to arrange for us to receive the referral prior to the procedure.

Additional information can be found at our website www.mwgi.com.

You may receive THREE separate bills for this procedure: 1 - Physician, 2 - Facility, 3 – Pathology

___ Colonoscopy ___ Flex Sig ___ EGD/Colon ___ EGD/Flex Sig

YOUR PROCEDURE IS SCHEDULED AT THE FOLLOWING TIME AND PLACE:

- Mountain West Endoscopy Center, 6360 South 3000 East, Ste 320, SLC, UT.**
801-944-3166
If not contacted by the day before your procedure, call 801-944-3166 to pre-register.
- Intermountain Medical Center, 5121 South Cottonwood Drive, Murray, UT.**
801-507-3262
Check in at the Eccles Outpatient Building, Level 2. Free Valet Parking Available.
Call the day before your procedure to pre-register, 801-314-4395
- Davis Hospital & Medical Center, 1600 Antelope Drive, Layton, UT.**
801-807-7034
Use the ER Entrance
- Lakeview Hospital, 630 E Medical Dr, Bountiful, UT.**
801-299-4657
South Entrance, Same Day Surgery Registration.
- Lakeview Endoscopy Center, 620 E Medical Dr, Suite 200, Bountiful, UT.**
801-299-6760
Call the day before your procedure to pre-register
- LDS Hospital, 8th Avenue & C Street, SLC, UT. ~Dr. Batista Only~**
801-408-1057
Entrance West Wing of Hospital. Call the day before your procedure to pre-register, 801-314-4395. Free Valet Parking Available.
- Riverton Hospital, 3741 West 12600 South, Riverton, UT.**
801-285-2775
Main entrance (Northeast corner) sign in at kiosk, take a pager and admitting will come get you from the waiting area.

FOR QUESTIONS PLEASE CALL OUR OFFICE AT _____