



Mountain West Gastroenterology

Colonoscopy Prep Instructions – OSMO Prep

<input type="checkbox"/> Oscar A. Batista, M.D.	<input type="checkbox"/> Edward J. Frech, M.D.	<input type="checkbox"/> Randall J. Ryser, M.D.
<input type="checkbox"/> Mark E. Boschert, M.D.	<input type="checkbox"/> Gordon E. Harmston, M.D.	<input type="checkbox"/> Michael J. Sossenheimer, M.D.
<input type="checkbox"/> Christopher C. Canale, M.D.	<input type="checkbox"/> Clarke A. Hilbig, M.D.	<input type="checkbox"/> I. Raymond Thomason, M.D.
<input type="checkbox"/> Brett W. Doxey, M.D.	<input type="checkbox"/> Robert G. Jones, M.D.	
<input type="checkbox"/> Peter C. Fenton, M.D.	<input type="checkbox"/> D. Rayburn Moore, M.D.	

7 DAYS PRIOR TO YOUR EXAM:

- If you have a heart condition, had a recent heart attack, stroke or cardiac stent placement, are taking **Aspirin and/or Plavix**, you should discuss this with your Cardiologist and/or Primary Care Physician prior to stopping any of these anti-clotting agents. While aspirin alone likely does not increase bleeding risks, the combination of Aspirin and Plavix may well increase the procedure risk. Yet stopping both or one of these agents may not be safe without discussing this with your Cardiologist and/or Primary Care Physician.
- It is equally important to discuss and review the usage of **Coumadin**. An appropriate taper or substitution may be needed. You should discuss this with your Cardiologist, Neurologist or Primary Care Physician and let our office know how to manage your blood thinners prior to the procedure.
- If you have **diabetes**, please follow our attached guidelines and discuss this with your Primary Care Physician.

5 DAYS PRIOR TO YOUR EXAM:

Avoid eating high fiber foods, fiber supplements and discontinue any **iron** supplements 5 days prior to your examination.

3 DAYS PRIOR TO YOUR EXAM:

If possible avoid taking anti-inflammatory medicines such as **Ibuprofen, Advil, Naproxen, Excedrin, Aleve or similar**. You can take Tylenol for pain.

Have your prescription for OSMO Prep Tablets filled at your pharmacy.

2 DAYS PRIOR TO YOUR EXAM:

You may eat a regular dinner, but stop eating solids after **7:00 pm**. Thereafter you may continue to drink clear liquids only. This means liquids you can see through such as Apple or White Grape juice, Water, Cola, Sprite, Jell-O, Coffee (No Creamer), Tea, Chicken or Beef broth, Gatorade, PowerAde, or Crystal Light. ***NO milk or pulp products: do not drink anything with red, purple or orange dyes.**

1 DAY PRIOR TO YOUR EXAM: *You may not eat any solid food until after your procedure!*

One of the most important aspects to a successful prep is to consume plenty of fluids the day before the exam.

Restrict your diet to clear liquids only, no solids. Liquids can be taken without restriction.

At 6:00 pm begin taking 4 tablets with a full 8 oz glass of any clear liquid every 15 minutes for a total of 5 doses (20 tablets). Make a conscious effort to drink plenty of clear liquids before, during, and after taking your Osmo Prep. Drink at least 24 ounces of liquids before going to bed. You may drink liquids throughout the night and up to 2 hours prior to your procedure. Your stool should be clear, yellow or green before your procedure.

THE DAY OF YOUR EXAM: *You may not eat any solid food until after your procedure!*

- If you have diabetes, test your blood sugar before coming in for your exam and bring the reading with you. If you take oral medications for your diabetes, don't take them the morning of your procedure unless otherwise instructed by your doctor. If you take insulin, bring your insulin with you. If you take insulin in the morning, take only ½ of your usual dose.
- Remember to **TAKE YOUR USUAL MEDICATIONS** (no blood thinners, no NSAIDS, no aspirin, no Plavix) the morning of your procedure.
- **5 hours before your scheduled arrival time at the endoscopy center**, begin taking 4 tablets every 15 minutes for a total of 3 doses (12 tablets). You may drink clear liquids up to 2 hours before your

scheduled arrival time. **Please have nothing by mouth 2 hours prior to your arrival, including water, candy or gum. Having anything by mouth within 2 hours before the procedure will result in postponing or cancelling your procedure.**

- **Nursing Mothers** – please “pump and dump” twice after the procedure before resuming breastfeeding.

You may apply zinc oxide, aloe wipes or petroleum jelly to anal area to prevent irritation. Baby Wipes or Medicated Tucks may be considered. Please bring someone with you to the endoscopy center to drive you home, as you will be sedated/anesthetized for the exam. Taxi drivers are not allowed. You may not drive for 12 hours. Your exam will be cancelled or must be done without sedation if you do not have someone to drive you home. You will be in the Endoscopy Lab for approximately two to three hours. The doctor will talk to you after the exam and will give you recommendations for diet, medication, follow up care, etc. Wear comfortable clothing; bring your glasses, hearing aids, insurance card(s) and completed information form. For your safety, no jewelry should be worn anywhere on your body the day of your procedure. No valuables should be brought to the center. We are not responsible for patient valuables. We will expect payment of co-pays, coinsurance and deductibles at the time of service. If your insurance requires a referral, you should contact your primary care physician to arrange for us to receive the referral prior to the procedure.

Additional information can be found at our website www.mwgi.com.

You may receive FOUR separate bills for this procedure: 1 - Physician, 2 - Facility, 3 – Pathology, 4-Anesthesia

___ Colonoscopy ___ Flex Sig ___ EGD/Colon ___ EGD/Flex Sig

YOUR PROCEDURE IS SCHEDULED AT THE FOLLOWING TIME AND PLACE:

- Mountain West Endoscopy Center, Old Mill Med Bldg., 6360 South 3000 East, Ste 320, SLC, UT 801-944-3166**
If not contacted by the day before your procedure, call to pre-register.
- Intermountain Medical Center, 5121 South Cottonwood Drive, Murray, UT 801-507-3262**
Check in at the Eccles Outpatient Building, Level 2. Free Valet Parking Available.
Call the day before your procedure to pre-register, 801-442-8600 or toll free 888-269-8674
- Davis Hospital & Medical Center, 1600 Antelope Drive, Layton, UT.**
801-807-7034 Use the ER Entrance
- Lakeview Hospital, 630 E Medical Dr, Bountiful, UT.**
801-299-4657
South Entrance, Same Day Surgery Registration.
- Lakeview Endoscopy Center, 620 E Medical Dr, Suite 200, Bountiful, UT.**
801-299-6760
Call the day before your procedure to pre-register
- LDS Hospital, 8th Avenue & C Street, SLC, UT. ~Dr. Batista Only~ 801-408-1057**
Entrance West Wing of Hospital. Call the day before your procedure to pre-register, 801-442-8600 or toll free 888-269-8674. Free Valet Parking Available.
- Riverton Hospital, 3741 West 12600 South, Riverton, UT.**
Riverton Hospital will call you prior to your procedure.

For Questions please call one of our offices:

Bountiful @ 801-298-0057
Old Mill @ 801-944-3141

Murray/TOSH @ 801-424-6310
Riverton @ 801-727-4280